


Folate and Arsenic Metabolism: A double -blind placebo controlled folate supplementation trial in Bangladesh.

J. Richard Pilsner, MPH, MPhil


Dr. Mary V. Gamble's Laboratory
Mailman School of Public Health
Columbia University

RO1 ES011601 & P42 ES10349

Arsenic metabolism by mono- and dimethylation.



Overview of One-Carbon Metabolism



Background: Arsenic Methylation & Folate

• Animal studies

- Dietary folate deficiency
 - *Toxicol.Lett. 2003: 145:167-74*

- Dietary methyl donor deficiency
 - *Mutat Res 1997: 386:315-34*

**Decrease total
urinary arsenic
excretion,
particularly DMA**

• Case-control studies in Taiwan

- Lower DMA in urine increased the risk of skin and bladder cancers and peripheral vascular disease

J Occup environ Med 2003, 45:241-8
Cancer Epidemiol Biomarker Prev 1997: 589-96
Toxicol Appl Pharmacol 2005, 206: 299-308

Background: Arsenic Methylation in Bangladesh

- Prospective cohort
 - Increase %MMA in urine is a risk factor for skin lesions
- Cross-Sectional Study

	%InAs	%MMA	%DMA
Folate (nM)	-0.12*	-0.12*	0.14*
Homocysteine (μM)	0.06	0.21 [#]	-0.14**

Spearman correlation coefficients * $p < 0.05$, ** $p < 0.001$, [#] $p < 0.001$

Environ Health Perspect 2005, 113:1683-88

Placebo-controlled Folate Intervention Trial

Hypothesis


Folate supplementation to folate deficient
Bangladeshi adults enhances
the methylation of arsenic.

Study Design:

Placebo-controlled Folate Intervention Trial


- 200 folate-deficient Bangladeshi adults
 - (Plasma folate < 9 nM)
- 12 weeks: Folic Acid (400 µg/d) or placebo
- Urinary As metabolites analyzed:
 - 0, 1, & 12 weeks
- All participants received a supply of multivitamins upon completion of the study

Results: Nutritional Parameters Pre- and Post-Intervention




a: $p < 0.0001$ within group; b: $p < 0.0001$ between groups; c: $p < 0.001$ between groups
Wilcoxon rank sum test for continuous variables

Results: Effects of Folate Supplementation on Arsenic Metabolites in Urine



*p < 0.05; ** p < 0.001 (folate vs. placebo); # p ≤ 0.05 (placebo pre- vs. post intervention)
Repeated measures linear regression

Distribution of %DMA in Urine after 12 Weeks Folate or Placebo Supplementation



- Summary:

- Folate intervention study indicates a causal relationship between folate supplementation and arsenic methylation in a folate-deficient population in Bangladesh.

- Implications:

- Enhancing arsenic methylation, as achieved by adequate folate status, could reduce arsenic-induced health outcomes.

- Future research:

- In a nested case-control study, we will determine if folate deficiency is a risk factor for subsequent development of arsenic-induced skin lesions.

Special Thanks...

Mary V. Gamble

Vesna Ilievski
Merle Jalakas

Epidemiology

Habibul Ahsan
Yu Chen
Pamela Factor-Litvak

Bangladesh

Tarique Islam
Shafiul Alam
Mominul Islam
Field Staff

Joseph Graziano

Vesna Slavkovich
Marni Hall
Olga Balac
Michael Jerome

Environmental Health Sciences

Faruque Parvez

Biostatistics

Xinhua Liu
Diane Levy

Lamont

Lex van Geen